Original Research Article

Study on assessment of Obstructive sleep apnea (OSA) risk in Obese pregnant women

Durga K¹, S Yuvarajan², Praveen Radhakrishnan^{2,*}, Antonious Maria Selvam²,
Arun Selvaraj³, Vinod Krishnadas³

¹Dept. of Obstetrics and Gynaecology, Sri Lakshmi Narayana Institute of Medical Science Medical College & Hospital, Kudapakkam, Puducherry, India

²Dept. of Respiratory Medicine, Sri Manakula Vinayagar Medical College, Kalitheerthalkuppam, India

³Dept. of Psychiatry, Sri Manakula Vinayagar Medical College, Kalitheerthalkuppam, India

ARTICLE INFO

Article history:

Received 16-06-2020

Accepted 09-07-2020

Available online 16-09-2020

Keywords:

OSA

Pregnancy

Obesity

STOP BANG

ABSTRACT

Introduction: Obstructive sleep apnea is a major public health concern which is associated with cardiovascular and cerebrovascular comorbidities. Eventhough association of OSA with adverse fetomaternal outcomes is documented, screening for OSA in pregnancy is a challenging task. So the purpose of our study is to assess the risk of OSA in obese pregnant ladies by using established STOP BANG Questionnaire for OSA.

Materials and Methods: This cross sectional study on assessment of obstructive sleep apnea risk among the obese pregnant women was done in tertiary care centre in the rural population of Puducherry.

Results: 21 obese pregnant women were included in our study. The mean age of our study population was 29.5 years. Most of the study subjects (85.7%) had history of snoring both reported by them and their bed partners. History of tiredness was reported by 14 patients (66.6%). Choking or history of gasping was reported by only 4 patients (19%). Results of our study showed that 42.9% of the obese pregnant women had high risk, 42.9% had intermediate risk and 14.3% for OSA as per the STOP BANG scoring.

Conclusion: Thus our study subjects were having higher risk of OSA which needed further evaluation with polysomnography. Thus the STOPBANG questionnaire can be used to screen for obstructive sleep apnea (OSA) in obese women during pregnancy.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

Obstructive sleep apnea (OSA) may affect nearly 20 percent of obese pregnant women.¹ Hypertensive disorders of pregnancy, Gestational diabetes mellitus and cardiomyopathy are the major comorbidities associated with OSA in pregnancy. It has been well documented that OSA in pregnancy have 2-fold increased risk of developing preeclampsia and 1.5-3.5-fold increased risk of the developing of gestational diabetes mellitus.²

Obstructive sleep apnea is a major public health concern which increases the cardiovascular and cerebrovascular comorbidities. OSA screening is mandatory for specific

population which includes obese people, patients with Resistant /poorly controlled diabetes mellitus/hypertension and Chronic kidney diseases. Even though lot of pathophysiological factors increases the risk of OSA in pregnancy, available literature in these group of population is sparse.

Adverse fetomaternal outcomes were well documented in patients with OSA in pregnancy. Hence it is mandatory to screen these patients which is a challenging task. Polysomnography is the gold standard investigation for the diagnosis of OSA but limited by its cost, waiting period in health care setting with significant patient load and time consumption. Unattended level 3 home based sleep studies is emerging as reliable cost-effective method for diagnosing

* Corresponding author.

E-mail address: drpraveen@gmail.com (P. Radhakrishnan).

OSA in high clinical suspicions.

The snoring, tiredness, observed apnea, high Blood pressure, Body mass index(BMI), age, neck circumference, and male gender (STOP-Bang) questionnaire was specifically developed to meet the need for a reliable, concise, and easy-to-use screening tool for OSA. It consists of eight dichotomous (yes/no) items related to the clinical features of sleep apnea. The total score ranges from 0 to 8.

So the purpose of our study was to assess the prevalence of risks of OSA in obese pregnant ladies by using established STOP BANG Questionnaire.

2. Aims and Objectives

To study the risk of OSA in Obese Pregnant women using STOP BANG questionnaire.

3. Materials and Methods

3.1. Study Area

Antenatal Clinic, Dept. of OBG, SLIMS, Puducherry.

3.2. Study design

Institutional Cross-sectional study.

Study duration

Jan 2019 to Jan 2020.

3.3. Sample size

Sample size was calculated based on the prevalence and study subjects from the previous study using free cal software.³

3.4. Inclusion criteria

Obese pregnant women with BMI more than 30(Both primigravida and multigravida).

3.5. Exclusion criteria

1. Pregnant women with Thyroid dysfunction.
2. Pregnant women with craniofacial abnormalities.
3. Pregnant women with Upper airway pathology.
4. Patients who are unwilling to participate in our study.

21 Obese pregnant women were recruited in our present study after fulfilling inclusion criteria and exclusion criteria. The study was conducted after approval from institutional ethical committee. Written informed consent was taken from all of them. After collecting demographic details of each patient, Patient was examined. Neck circumference in centimeters /inches and BMI(weight/height in meter square) were recorded .Relevant investigation were sent to apply execution criteria. Risk of OSA was assessed using STOP BANG questionnaire. All the data was entered and analysed with epi info software version 3.4.3.

3.6. Risks assessment based on STOPBANG scoring

3.6.1. Low risk of OSA

Score of 0-2

3.6.2. Intermediate risk of OSA

Score of 3-4

3.6.3. High risk of OSA

Score of 5-8

or Yes to 2 or more of 4 STOP questions + male gender

or Yes to 2 or more of 4 STOP questions + BMI > 35 kg/m²

or Yes to 2 or more of 4 STOP questions + neck circumference

4. Results

21 obese pregnant women were included in our study. The mean age of our study population was 29.5years. All of them were from rural areas in the region of Puducherry.70 % of the study population belong to poor socioeconomic status as per Kuppusamy scale.

Out of 21 obese pregnant women, 66%(n=14) were multigravida and 34% were primigravida.10 % of them had history of previous abortions.

5 (23.8%) of them had Pregnancy induced hypertension 7(33.3%) had gestational diabetes.

Regarding symptoms pertaining to OSA, most of them had snoring (n=18, 85.7%) followed by day time sleepiness (n=15,71.4%), Day time tiredness (n=14, 66.6%) Early morning headache (n=6, 28.5%) and choking episodes (n=4,19%), and nocturnal enuresis in 2 patients.

66.6%(n=14)of the study population had BMI in the range 30-35 whereas 28.6%(n=6) had BMI between 35 to 40. one patient had morbid obesity with BMI more than 40.

With regard to stop bang questionnaire, Most of the study subjects (85.7%) had history of snoring both reported by them and their bed partners. History of tiredness was reported by 14 patients (66.6%). choking or history of gasping was reported by only 4 patients (19%). 4 of them had systemic hypertension and 5 patients had pregnancy induced hypertension.

13(61.9%) patients had BMI greater than 35. Question pertaining to age and sex in questionnaire is not applicable for our study population.

Table 1: Distribution of Study subjects based on Body mass index

BMI	n(%)
30-35	14(66.6)
35-40	6(28.6)
40	1(4.8)

Table 2: Scoring of stop bang questionnaire

Stop	Yes N(%)	No. N(%)
SNORING (S) Do you Snore Loudly ?	18(85.7)	3(14.3)
TIREDNESS?(T)	14(66.6)	7(33.3)
Observed ?(O) Has anyone noticed Choking episodes?	4(19)	17(80.9)
Pressure ?(P) History of High Blood Pressure ?	9(42.8)	12(57.1)
Body Mass Index more than 35 kg/m2?(B)	7(33.3)	14(66.7)
Age older than 50 ? (A)	Not Applicable	Not applicable
Neck size large ? (N) For male, 17 inches / 43cm or larger? For female, 16 inches / 41cm or larger?	12(57.1)	9(42.9)
Gender = Male ? (G)	Not applicable	Not applicable

Fig. 1: Distribution of STOP BANG in study subjects**Table 3:** Risk of OSA based on stop bang score

OSA Risk	n(%)
Low risk	3(14.3)
Intermediate risk	9(42.9)
High risk	9(42.9)

5. Discussion

This cross sectional study on assessment of obstructive sleep apnea risk among the obese pregnant women was done in tertiary care centre in the rural population of Puducherry.

Results of our study showed that 42.9% of the obese pregnant women had high risk, 42.9% had intermediate risk and 14.3% for OSA as per the STOP BANG scoring. Thus most of our study subjects were having higher risk of OSA which needed further evaluation with polysomnography.

Chung et al⁴ showed STOP-BANG with specific combinations of individual components in the questionnaire improved the specificity. Patients with a STOP score of ≥ 2 , male gender and a BMI greater than 35 kg/m² were

more predictive of OSA than age greater or equal to 50 and neck circumference > 40 cm. So, a screening tool is much needed which should be specific for the women in pregnancy to diagnose OSA.

Haulek Mergan et al⁵ in their study with 217 obese patients, scoring cut off 3-4 provides a better balance of sensitivity and specificity for all OSA.

9 patients had hypertension in our study population among the obese pregnant women. Previous studies showed high incidence of Ischemic heart disease, systemic hypertension, stroke, Acid peptic disease, heart failure with OSA.^{6,7}

Jennifer E. Dominguez et al⁸ in their study noted that higher total STOP-BANG score was clearly linked with high likelihood of pre-pregnancy and pregnancy-induced hypertension with diabetes mellitus which is similar to our study. 42.8% of our study subjects were hypertensive with higher STOP BANG scores.

STOP-BANG was the most commonly used screening tool among the departments that screened for OSA.⁹

Similar studies with the use of STOPBANG in the first and third trimesters of pregnancy concluded that even though it had a good negative predictive value, the score was not useful for detecting OSA.⁹⁻¹³ Tantrakul et al¹⁴ studied the usage of STOPBANG questionnaire in 72 women across trimesters and found it to have the best predictive value during the second trimester.

In our study, 12 patients (57.1%) had neck circumference more than 16 inches. In a study by Fiona Pearson et al¹⁵ 41% of study patients had neck circumference greater than 40 cm.

Most of the obese pregnant women (87.5%) in our study reported snoring. Other authors have described snoring rates of 17-34% in pregnancy but not in obese pregnant women^{16,17}. Snoring was prevalent in our study this may be due to the fact that other similar studies concentrated on snoring conducted detailed interviews with precise criteria for snoring, whereas our study used simple self-reporting with the help STOP BANG questionnaire. Smith et al¹⁸ suggested that habitual snoring (snoring at 14 least 3 nights per week) could be used as a triage tool for OSA in pregnancy, especially among those with a BMI ≥ 40 .

A large proportion of study population reported day time sleepiness (71.4%) and tiredness (66.6%) which is similar to a study by Fiona Pearson et al.¹⁵ Patients also had history of choking episodes which is predominantly seen in patients with BMI greater than 35. Few patients reported early morning headache and nocturnal enuresis.

Thus patients with higher BMI, neck circumference more than 16 inches with higher reporting of snoring and tiredness invariably had higher STOP BANG scores which shows these group of patients have high risk of developing obstructive sleep apnea in pregnancy like in general population. However these patients should be

referred for polysomnography to confirm the diagnosis of Obstructive sleep apnea.

6. Conclusion

This study showed higher prevalence of risks for OSA in obese pregnant women as per STOP BANG questionnaire which can be utilised for screening OSA in these group of patients. However it requires meticulous history taking and scoring with the questionnaire which in turn requires measurements that are not normally done in an antenatal outpatient department. Scoring of STOP BANG questionnaire is much simpler and less time consuming when compared to other questionnaires to assess the risk of OSA. Thus Obese pregnant women reporting loud and frequent snoring with day time tiredness and comorbidities should be referred for investigation of obstructive sleep apnea. However, Multicentric studies with large sample size should be initiated using Level 1 or Level 2 Sleep study in obese women with the aim of developing a pregnancy-specific screening tool for OSA.

7. Acknowledgement

I acknowledge my post graduates Dr. Vinayak Nandhanan, Dr. Prathaban, Dr. Madhan, Dr. Ijas for their timely help.

8. Source of Funding

None.

9. Conflict of Interest

None.

References

- Pien GW, Pack AI, Jackson N, Maislin G, Macones GA, Schwab RJ, et al. Risk factors for sleep-disordered breathing in pregnancy. *Thorax*. 2014;69(4):371–7.
- Facco FL, Parker CB, Reddy UM, Silver RM, Koch MA, Louis JM, et al. Association Between Sleep-Disordered Breathing and Hypertensive Disorders of Pregnancy and Gestational Diabetes Mellitus. *Obstet Gynecol*. 2017;129(1):31–41.
- El-Helbawy R, Elmahlawy I, Shaheen A, Reda A. Ibrahim c Obstructive sleep apnea in pregnancy. Is it a new syndrome in obstetrics?, . *Egypt J Chest Dis Tuberc*. 2017;66(4).
- Chung F, Yang Y, Brown R, Liao P. Alternative Scoring Models of STOP-Bang Questionnaire Improve Specificity To Detect Undiagnosed Obstructive Sleep Apnea. *J Clin Sleep Med*. 2014;10(9):951–8.
- Mergen H, Altındağ B, Uçar ZZ, Kılıçaslan IK. The Predictive Performance of the STOP-Bang Questionnaire in Obstructive Sleep Apnea Screening of Obese Population at Sleep Clinical Setting. *Cureus*. 2019;11(12):6498.
- Dincer FF, Neil WO. Deleterious effects of sleep-disordered breathing on the heart and vascular system. *Respir*. 2006;73:124–30.
- Ing AJ, Ngu MC, Breslin ABX. Obstructive sleep apnea and gastroesophageal reflux. *Am J Med*. 2000;108(4):120–5.
- Dominguez JE, Grotegut CA, Cooter M, Krystal AD, Habib AS. Screening extremely obese pregnant women for obstructive sleep apnea. *Am J Obstet Gynecol*. 2018;219(6):613.e1–e10.
- Miskovic A, Simpson M, Dominguez JE, Bullough AS. Recognising obstructive sleep apnoea in pregnancy - a survey of UK obstetric anaesthetists. *Anaesth*. 2017;72(12):1564.
- Lockhart EM, Abdallah AB, Tuuli MG, Leighton BL. Obstructive Sleep Apnea in Pregnancy. *Obstet Gynecol*. 2015;126(1):93–102.
- Facco FL, Ouyang DW, Zee PC, Grobman WA. Development of a Pregnancy-Specific Screening Tool for Sleep Apnea. *J Clin Sleep Med*. 2012;08(04):389–4.
- Nagappa M, Liao P, Wong J, Auckley D, Ramachandran SK, Memtsoudis S, et al. Validation of the STOP-Bang Questionnaire as a Screening Tool for Obstructive Sleep Apnea among Different Populations: A Systematic Review and Meta-Analysis. *PLOS ONE*. 2015;10(12):e0143697.
- Sequeira T, Bublit M, Adodoadji E, Livingston Z, Bourjeily G. Stopbang Questionnaire Correctly Detects the Absence of Obstructive Sleep Apnea in the First Trimester of Pregnancy. *Sleep*. 2017;40(1):A186.
- Tantrakul V, Sirijanchune P, Panburana P, Pengjam J, Suwansathit W, Boonsarngsuk V, et al. Christian Guilleminault Screening of Obstructive Sleep Apnea during Pregnancy: Differences in Predictive Values of Questionnaires across Trimesters. *J Clin Sleep Med*. 2015;11(2):157–63.
- Pearson F, Batterham AM. Sean Cope The STOPBANG Questionnaire as a Screening Tool for Obstructive Sleep Apnoea in Pregnancy. *J Clin Sleep Med*. 2019;15(5):705–10.
- Ge X, Tao F, Huang K, Mao L, Huang S, Niu Y, et al. Erigene Rutayisire Maternal Snoring May Predict Adverse Pregnancy Outcomes: A Cohort Study in China. *PLoS One*. 2016;11(2).
- O'Brien LM, Bullough AS, Owusu JT, Tremblay KA, Brincat CA, Chames MC, et al. Pregnancy-onset habitual snoring, gestational hypertension, and preeclampsia: prospective cohort study. *Am J Obstet Gynecol*. 2012;207(6):487.e1–e9.
- Smith J, Triebwasser JE, Langen E, Brien O, M L. How Best to Screen for Obstructive Sleep Apnoea in Pregnancy? *AJOG*. 2018;218(1).

Author biography

Durga K Associate professor

S Yuvarajan Professor

Praveen Radhakrishnan Assistant Professor

Antonious Maria Selvam Assistant Professor

Arun Selvaraj Professor

Vinod Krishnadas Associate Professor

Cite this article: Durga K, Yuvarajan S, Radhakrishnan P, Selvam AM, Selvaraj A, Krishnadas V. **Study on assessment of Obstructive sleep apnea (OSA) risk in Obese pregnant women.** *IP Indian J Immunol Respir Med* 2020;5(3):181–184.