Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Immunology and Respiratory Medicine

Journal homepage: <https://www.ijirm.org/>

Case Report

Acute pulmonary embolism after short term exposure to smoke in an enclosed area

Sayan Malakar¹, Bhagwan Dass Negi¹, Katyayani Dutt¹, Krishna Bharath¹,
 Sujeet Raina^{1,*}

¹Dept. of Medicine, Dr. Rajendra Prasad Govt. Medical College & Hospital, Tanda, Kangra, India

ARTICLE INFO

Article history:

Received 09-12-2021

Accepted 27-12-2021

Available online 31-12-2021

Keywords:

Embolism

Inflammation

Smoke

ABSTRACT

We report a case of acute pulmonary embolism (PE) following short term exposure to smoke in an enclosed area. The patient was obese and had type 2 diabetes mellitus. He developed PE as a consequence of acute systemic inflammatory response to short term exposure to smoke and an underlying chronic inflammatory milieu.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](#), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Pulmonary thromboembolism is known among one of “the Great Masquerader” with a grave prognosis if not diagnosed and managed early. High index of suspicion is required for making the diagnosis as the symptoms and signs are nonspecific. A high Wells score has a higher probability of PE.¹ The overall risk of venous thromboembolism (VTE) depends on the predisposing environmental and genetic factors. VTE follows as consequence of the interaction between patient related usually permanent risk factors and setting-related usually temporary risk factors. The risk increases in proportion to the number of predisposing factors.² Diabetes and obesity are considered to be weak risk factors with an odds ratio <2.² Here we are reporting a case who presented with sudden onset shortness of breath after short term exposure to smoke in an enclosed area and on investigations was diagnosed to have acute PE. This case is presented for the following reasons. Firstly, among the common predisposing factors for VTE smoke inhalation as a provoking factor is rarely described. Acute PE after short term exposure to smoke in an enclosed area

is not frequently reported. Secondly, the hs-CRP at the admission was higher in comparison to a followup level. The inflammatory marker levels suggest that short term exposure to smoke in an enclosed area acted as a trigger and initiated acute systemic inflammation in an environment of chronic inflammatory state. Thirdly, the patient had no evidence of inhalational lung injury as a result of exposure to smoke or chemical products of combustion. This patient had no upper airway obstruction, atelectasis, pulmonary oedema and infection or chemical pneumonitis. Fourthly, The clinical probability for PE was low on three level score and unlikely on two level score of Wells point score.¹ Lastly, an episode of syncope as one of the presenting symptom of PE.

2. Case Report

A 45-year-old male presented with history of dyspnoea which was sudden in onset and progressive for the last 36 hours. Patient was apparently well before the onset of symptoms. On the pre Diwali eve while returning home from the market he noticed smoke coming from his office room located on the ground floor of his house. He opened the door of the room and immediately inhaled the gush of

* Corresponding author.

E-mail address: sujeetrashmishera@yahoo.co.in (S. Raina).

smoke. He entered smoke filled room and found a mattress burning. He switched off all the electrical appliances and rushed out. He didn't suffer any burns. Two hours later he developed dyspnoea on exertion and consulted the local community health centre. He was reassured and sent back home. The patient continued to have dyspnoea on exertion as well as on rest throughout the night, the next day which was the Diwali day as well as on the Diwali night. In the morning after the festival while walking in his courtyard he had an episode of loss of consciousness for approximately two minutes. He recovered spontaneously and was rushed to the hospital. There were no history of fever, chest pain, palpitations, wheeze, cough, haemoptysis, stridor, dysphonia, dysphagia and sialorrhoea. He was a diabetic for the last four years with poor diabetes control, non-smoker and consumed alcohol on social occasions. He had no significant past history. On examination his blood pressure was 76/48 mm of Hg; pulse rate was 112/minute, respiratory rate was 34/minute and had cool sweaty periphery with prolonged capillary refill time. All peripheral pulses were palpable without any delay. SaO₂ was 86% on ambient air which improved after oxygenation at the rate of 4-5 L / minute with a facemask. GCS was 15 / 15 without any other focal neurological signs. Jugular venous pressure was not raised. Systemic examination of the respiratory and cardiovascular systems was normal. Review of other systems was normal. His electrocardiogram revealed sinus tachycardia, QTc of 443 msec, left anterior hemiblock and poor progression of R wave from V1-V6 (Figure 1). Central venous pressure (CVP) was 19 cm of water. Patient was started on inotrope infusion. On investigations total leucocyte count was 11,500/cmm. Rest of the hemogram, serum electrolytes, renal, liver functions and lipid profile were normal. Random blood glucose was 335mg/dl and HbA1c was 12.7%. His high sensitivity cardiac troponin I (hs-cTnI) on admission was 23.5 pg/ml [range 28.9-39.0]. The brain natriuretic peptide (BNP) was 334pg/ml. D-dimer was raised and semi-quantitative determination of plasma d-dimer by slide agglutination was >1.0 < 2.0(normal <0.5). His high sensitivity C-reactive protein (hs-CRP) was 25.4 mg/L. Chest x-ray was normal. His bedside echocardiogram revealed mild tricuspid regurgitation (TR) and pulmonary artery hypertension (PAH). Computed tomography (CT) pulmonary angiography revealed massive thrombus in left pulmonary artery extending into segmental branches (Figure 2). Patient was managed with Streptokinase 0.25 million IU stat over 30 minutes followed by 0.1 million IU/h for next 24 hours. He improved hemodynamically and inotrope was tapered off within 12 hours. Compression venous ultrasound of both lower limbs was normal. His BMI was 32kg/m². After initial five days of low molecular weight Heparin patient was shifted to oral Dabigartan and discharged. His follow up hs-CRP 10 days later was 6.8 mg/L. His follow up echocardiogram after two months revealed

mild PAH and mild TR. An informed consent of the patient has been taken for publication of this data and images before submission of the manuscript.

Fig. 1: ECG showing sinus tachycardia,QTc of 433 msecs, left anterior hemiblock, poor progression of R wave from V1-V6

Fig. 2: CT pulmonary angiography axial section showing filling defect in left pulmonary artery (arrow) also extending into segmental branches

3. Discussion

Venous thromboembolism (DVT and PE) is not uncommon in India as traditionally believed. The Indian population also have similar predisposing (risk) factors as their counterparts in the western world and there appears no reason to believe that the prevalence should be any different in India.³ Traditional predisposing factors for VTE have been described as strong, moderate and weak with specific odds ratios for each group.² The patient under discussion had none of the strong or moderate

risk factors but had diabetes mellitus and obesity among weak risk factors. We presume that the presence of the two weak risk factors and additional exposure to smoke in a closed space resulted in acute pulmonary embolism. The evidence in favour of the presumption is that patient had a higher hs-CRP level at admission as compared to follow up thus explaining an acute on chronic inflammatory state. Inflammation and thrombosis are interrelated. Inflammation is a trigger for endothelial dysfunction, platelet activation and hypercoagulability. A growing interest about air pollution as a risk factor of acute venous thromboembolism has been observed.^{4–6} Air pollution is a heterogeneous, complex mixture of gases, liquids, and particulate matter. Epidemiological studies have linked cardiovascular events in relation to both short and long term exposure to present day ambient air pollution. The biological mechanisms involved include enhanced coagulation/thrombosis, a propensity for arrhythmias, acute arterial vasoconstriction, systemic inflammatory responses, and the chronic promotion of atherosclerosis.⁷ In a retrospective case-control study one month's exposure to elevated levels of different air pollutants was identified as a possible risk factor for the development of acute isolated PE. It was concluded that the onset of acute isolated (i.e. without deep vein thrombosis) pulmonary embolism (PE) is due to a potential direct damage by exposure to fine particulate matter to the lung vasculature.⁸ To summarize, exposure to air pollution should be considered a risk factor for unprovoked acute isolated PE.

4. Acknowledgements

None.

5. Conflicts of interests

The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article

6. Source of Funding

The authors received no financial support for the research, authorship, and/or publication of this article

References

1. Wells PS, Anderson DR, Rodger M, Ginsberg JS, Kearon C, Gent M, et al. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer. *Thromb Haemost.* 2000;83(3):416–20.
2. Konstantinides SV, Torbicki A, Agnelli G, Danchin N, Fitzmaurice D, Galie N, et al. ESC guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J.* 2014;35(43):3033–69. doi:10.1093/eurheartj/ehu283.
3. Lee AD, Stephen E, Agarwal S, Premkumar P. Venous thromboembolism in India. *Eur J Vasc Endovasc Surg.* 2009;37(4):482–5. doi:10.1016/j.ejvs.2008.11.031.
4. Baccarelli A, Martinelli I, Zanobetti A, Grillo P, Hou LF, Bertazzi PA, et al. Exposure to particulate air pollution and risk of deep vein thrombosis. *Arch Intern Med.* 2008;168(9):920–7. doi:10.1001/archinte.168.9.920.
5. Baccarelli A, Martinelli I, Pegoraro V, Melly S, Grillo P, Zanobetti A, et al. Living near major traffic roads and risk of deep vein thrombosis. *Circulation.* 2009;119(24):3118–24. doi:10.1161/CIRCULATIONAHA.108.836163.
6. Dales RE, Cakmak S, Vidal CB. Air pollution and hospitalization for venous thromboembolic disease in Chile. *J Thromb Haemost.* 2010;8(4):669–74. doi:10.1111/j.1538-7836.2010.03760.x.
7. Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, et al. Air pollution and cardiovascular disease: a statement for healthcare professionals from the Expert Panel on Population and Prevention Science of the American Heart Association. *Circulation.* 2004;109(21):2655–71. doi:10.1161/01.CIR.0000128587.30041.C.
8. Spiezia L, Campello E, Bon M, Maggiolo S, Pelizzaro E, Simioni P, et al. Short-term exposure to high levels of air pollution as a risk factor for acute isolated pulmonary embolism. *Thromb Res.* 2014;134(2):259–63. doi:10.1016/j.thromres.2014.05.011.

Author biography

Sayan Malakar, Ex-Post Graduate Student

Bhagwan Dass Negi, Ex-Post Graduate Student

Katyayani Dutt, Ex-Post Graduate Student

Krishna Bharath, Ex-Post Graduate Student

Sujeet Raina, Associate Professor

Cite this article: Malakar S, Negi BD, Dutt K, Bharath K, Raina S. Acute pulmonary embolism after short term exposure to smoke in an enclosed area. *IP Indian J Immunol Respir Med* 2021;6(4):253-255.